

Optical Interferometry with Pulsed Fields

Robert W. Schoonover¹, Brynmor J. Davis¹, Randy A. Bartels², P. Scott Carney¹

¹ University of Illinois-Department of Electrical and Computer Engineering
Urbana, Illinois 61801

² Colorado State University-Department of Electrical and Computer Engineering
Boulder, CO 80309
Email: rschoono@uiuc.edu

Abstract: An analysis of coherence properties of pulsed fields in interferometric experiments is presented. The results bear on means to recover certain statistical properties of the source in a two-pinhole experiment.

©Optical Society of America

OCIS codes: (030.6600) Statistical Optics (120.3180) Interferometry

Statistical optics, or coherence theory, is the study of the outcome of measurements made with non-deterministic light. Until recently, almost all optical experiments could be well-modeled within the framework of stationary, at least in the wide sense, random processes. This is not the case with pulsed optical fields. Such non-stationary fields must be treated in a much more general setting.

The theoretical study of non-stationary statistical optics has seen a recent resurgence. In references [1, 2] intrinsically stationary sources [3], members of a restricted class of non-stationary random processes, are studied. A stationary field is modulated in a deterministic fashion. While this model is not the most general one to describe optical phenomena, it is a good bridge between a general theory of two-time, two-point measurements and standard coherence theory.

In this presentation, pulses are treated as resulting from the modulation of a stationary source. A mathematical model is formulated to relate observable quantities to the statistical properties of the source. Specifically, the intensity pattern formed in a two-pinhole experiment is considered for many different cases of non-stationary pulses. The duration of the pulse, the coherence time of the source, the time between pulses, and the integration time at the detector are all considered. Equivalent intensity profiles are shown for fields with different parameters. In certain cases, it is shown that the coherence properties of the unmodulated process can be recovered.

Simulations illustrate the role of detection in the relationship between observables and the underlying coherence properties of the field. Special attention is paid to cases where long-time-average measurements (in a certain sense) are unable to recover the basic statistical properties of the optical field.

References

- [1] H. Lajunen, J. Tervo, J. Turunen, P. Vahimaa, and F. Wyrowski, "Spectral coherence properties of temporally modulated stationary light sources," *Optics Express* **11**(16), 1894–1899 (2003).
- [2] H. Lajunen, J. Tervo, and P. Vahimaa, "Overall coherence and coherent-mode expansion of spectrally partially coherent plane-wave pulses," *Journal of the Optical Society of America A* **21**(11), 2117–2123 (2004).
- [3] R. Gase and M. Schubert, "On the Determination of Spectral Properties of Non-stationary Radiation," *Journal of Modern Optics* **29**(10), 1331–1347 (1982).